



endobioticum																	
Clavibacter michiganensis	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
<b>Potato spindle tuber viroid</b>	0	0	0	0	0	0	-	-	-	-	-	0	0	0	0	0	0
<b>Tomato stolbur</b>	0	0	0	0	0	0	-	-	-	-	-	-	-	-	-	-	-
<b>Meloidogyne chitwoodi and fallax</b>	0	0	0	0	0	0	-	-	-	-	-	0	0	0	0	0	0
<b>Ditylenchus destructor</b>	0	0	0	0	0	0	-	-	-	-	-	0	0	0	0	0	0
<b>3. Direct progeny tolerances</b>																	
Other varieties and off types	0	0.01	0.25	0.25	0.5	0.5	0.25 & 0.1	0.25 & 0.1	0.25 & 0.1	0.25 & 0.1	0.5 & 0.2	0	0.01	0.1 var 0.25 off	0.1 var 0.25 off	0.1 var 0.25 off	As class
Virus (%)	0	0.5	2 (1 severe)	4 (2 severe)	10 (5 severe)	10	4	4	4	4	10	0	0.5	4	4	4	As class

<sup>1</sup> Harmful organisms in bold are covered by the EU legislation on quarantine pests (Directive 2000/29/EC)

\* The figure in brackets is the allowable % surface area covered: a tuber is deemed to be affected by the disease only if surface area affected exceeds the specified allowable surface tolerance.

\*\* In Scotland there are 4 field generations permitted at PB; PB1, PB2, PB3 and PB4, and six field generations permitted at Basic SE1, SE2, SE3, E1, E2, E3, A is a fall grade only. In practice only 4 to 6 generations are normally used.